# Code No: 132ABR16JAWAHARLAL NEHRU TECHNOLOGICAL UNIVERSITY HYDERABADB. Tech I Year II Semester Examinations, August - 2018<br/>MATHEMATICS – II<br/>(Common to EEE, ECE, CSE, EIE, IT, ETM)

#### Time: 3 hours

**Note:** This question paper contains two parts A and B. Part A is compulsory which carries 25 marks. Answer all questions in Part A. Part B consists of 5 Units. Answer any one full question from each unit. Each question carries 10 marks and may have a, b, c as sub questions.

### PART-A

Find the Laplace transform of the function  $f(t) = t^2$ . 1.a) [2] Find Laplace transform of  $4\sin(t-3)$ . b) [3] Show that  $\Gamma(n) = 2 \int_0^\infty e^{-x^2} x^{2n-1} dx$ . c) [2] Show that  $\beta(p,q) = \check{\beta}(p+1,q) + \beta(p,q+1)$ . d) [3] Find the area bounded by the curves y = x,  $y = x^2$ . e) [2] Evaluate  $\int_{0}^{1} \int_{0}^{1} x^2 y^2 dx dy$ f) [3] If  $\phi = x^2 y^2 z^2$  then find Grad  $\phi$ . [2] g) Find a unit normal vector to the surface  $x^2 + y^2 + 2z^2 = 26$  at the point (2,2,3). [3] h) Find curl  $\overline{F}$  when  $\vec{F} = 3x^2i + (2xz - y)j + zk$ . i) [2]

- j) Is the work done by a force in moving a particle from one point to another point in an irrotational field is independent of the path of integration? Justify the answer. [3]
  - PART-B

## (50 Marks)

- 2. Use Laplace transforms, solve y''(t) + 5y'(t) + 6y(t) = t, y(0) = 1, y'(0) = 1. [10] OR
- 3. Solve by using Laplace transforms  $y'' + 4y' + 3y = e^{-t}$  with y(0) = y'(0) = 1. [10]

4. Prove that 
$$\int_0^1 \frac{x^2 dx}{\sqrt{1-x^4}} \times \int_0^1 \frac{dx}{\sqrt{1+x^4}} = \frac{\pi}{4\sqrt{2}}$$
 using  $\beta - \Gamma$  functions. [10]  
OR

5.a) Prove that 
$$\Gamma\left(\frac{1}{2}\right) = \sqrt{\pi}$$
  
b) Prove that  $\beta(m,n) = \frac{\Gamma(m) \Gamma(n)}{\Gamma(m+n)}$ . [5+5]

## WWW.MANARESULTS.CO.IN

Max. Marks: 75

(25 Marks)

The plane  $\frac{x}{a} + \frac{y}{b} + \frac{z}{c} = 1$  meets the axes in *A*, *B* and *C*. Find the volume of the tetrahedron 6. OABC. [10]

OR

7. Evaluate 
$$\int_{0}^{1} \int_{0}^{1-x^{-1}-x^{-y}} \int_{0}^{1-x^{-y}} x^2 yz \, dz \, dy \, dx$$
. [10]

,

Prove that if  $\vec{r}$  is the position vector of any point in space then  $r^n \vec{r}$  is irrotational and is 8. solenodial if n = -3. [10]

- Evaluate  $\nabla \cdot \left( r \nabla \left( \frac{1}{r^3} \right) \right)$  where  $r = \sqrt{x^2 + y^2 + z^2}$ . 9.a) If  $\overline{R} = x\overline{i} + y\overline{j} + z\overline{k}$ , then find  $\nabla .\overline{R}$  and  $\nabla \times \overline{R}$ . b) [5+5]
- Verify Stoke's theorem for the vector field  $\vec{F} = (x^2 y^2)i + 2xyj$  integrated round the 10. rectangle in the plane z = 0 and bounded by the lines x = 0, y = 0, x = a, y = b. [10]
  - OR
- Verify divergence theorem for  $2x^2yi y^2j + 4xz^2k$  taken over the region of first octant of the cylinder  $y^2 + z^2 = 9$  and x = 2. [10] 11.

--00000--

# WWW.MANARESULTS.CO.IN