

**S.E. ( Computer / Information Technology )**  
**Engineering Mathematics - III**  
**2012 Course**

Time: 2Hours

Max. Marks : 50

Instructions to the candidates:

- 1) Solve Q1 or Q2, Q3 or Q4, Q5 or Q6 and Q7 or Q8.
- 2) Neat diagrams must be drawn wherever necessary.
- 3) Figures to the right side indicate full marks.
- 4) Use of Calculator is allowed.
- 5) Assume Suitable data if necessary

Q1) a) Solve any **two** of the following [08]

i)  $(D^2 + 4) y = \cos 3x \cdot \cos x$

ii)  $(D^2 + 6D + 9) y = e^{3x} / x^2 \dots$  ( by variation of parameters method)

iii)  $x^3 (d^3y / dx^3) + 2x^2 (d^2y / dx^2) + 2y = 20 (x + 1/x)$

b) Obtain f(k) given that, [04]

$$f(k+2) + 5f(k+1) + 6f(k) = 0, \quad k \geq 0 \quad f(0) = 0, \quad f(1) = 2$$

by using Z transform.

**OR**Q2) a) An emf  $E \sin(pt)$  is applied at  $t = 0$  to a circuit containing a condenser 'C' and [04]

Inductance 'L' in series. The current 'x' satisfies the equation

$$L(dx/dt) + \frac{1}{C} \int x dt = E \sin(pt)$$

Where  $p = \frac{-dq}{dt}$ . If  $p^2 = \frac{1}{LC}$  and initially the current x and charge q is zero then show that current in the circuit at any time t is  $\frac{E}{2L} t \sin(pt)$ .

b) Solve the integral equation [04]

$$\int_0^{\infty} f(x) \cos \lambda x dx = \begin{cases} 1 - \lambda & 0 \leq \lambda \leq 1 \\ 0 & \lambda > 1 \end{cases}$$

And hence show that  $\int_0^{\infty} \frac{\sin^2 z}{z^2} dz = \pi/2$ c) Attempt **any one** [04]i) Find the Z- transform of  $f(k) = e^{-2k} \cos(5k + 3)$   
[www.manareresults.co.in](http://www.manareresults.co.in)

ii) Find the inverse Z- transform of  $\frac{z(z+1)}{z^2-2z+1}$ ,  $|z| > 1$

Q3) a) The first four moments of a distribution about 2 are 1, 2.5, 5.5 and 16. Calculate the first four moments about the mean, A. M., S. D.,  $\beta_1$  and  $\beta_2$  [05]

b) In a certain examination 200 students appeared. Average marks obtained were 50% with standard deviation 5%. How many students do you expect to obtain more than 60% of marks, supposing that the marks are distributed normally ? [04]  
( Given  $z=2$  ;  $A = 0.4772$  )

c) Find the directional derivatives of: [04]  
 $\phi = xy^2 + yz^2 + zx^2$  at (1,1,1) along the line  $2(x-2) = y+1 = z-1$

OR

Q4) a) Calculate the coefficient of correlation for the following data [05]

x	1	2	3	4	5	6	7	8	9
y	9	8	10	12	11	13	14	16	15

b) Prove the following (Any one) [04]

i)  $\nabla^4 r^4 = 120$

ii)  $\nabla \cdot \left[ r \nabla \frac{1}{r^5} \right] = \frac{15}{r^6}$

c) Show that  $\vec{F} = (x^2 - yz)\vec{i} + (y^2 - zx)\vec{j} + (z^2 - xy)\vec{k}$  is irrotational. Also find  $\Phi$  such that  $\vec{F} = \nabla\Phi$  [04]

Q5) a) Find the work done in moving a particle along [04]

$x=a \cos\theta, y= a \sin\theta, z=b\theta$ , from  $\theta = \frac{\pi}{4}$  to  $\theta = \frac{\pi}{2}$  under a field of force given

by  $\vec{F} = -3a \sin^2\theta \cos\theta \hat{i} + a(2\sin\theta - 3\sin^3\theta)\hat{j} + b \sin 2\theta \hat{k}$

b) Evaluate  $\iint_s (yz\hat{i} + zx\hat{j} + xy\hat{k}) \cdot d\vec{s}$ , where s is the curved surface of the cone [04]  
 $x^2 + y^2 = z^2, z = 4$

c) Using Stokes Theorem to evaluate  $\int_c (4y\hat{i} + 2z\hat{j} + 6y\hat{k}) \cdot d\vec{r}$  where C is the [04]  
curve of intersection of  $x^2 + y^2 + z^2 = 2z$  and  $x = z - 1$

OR

Q6) a) A vector field is given by  $\vec{F} = (2x - \cos y)\hat{i} + x(4 + \sin y)\hat{j}$ , evaluate  $\int_c \vec{F} \cdot d\vec{r}$ , [04]

where c is the ellipse  $\frac{x^2}{a^2} + \frac{y^2}{b^2} = 1, z = 0$ .

b) Prove that  $\iiint_v \frac{1}{r^2} dv = \iint_s \frac{1}{r^2} \bar{r} \cdot d\bar{s}$ , where  $s$  is closed surface enclosing the volume  $v$ . Hence evaluate  $\iint_s \frac{x\hat{i}+y\hat{j}+z\hat{k}}{r^2} \cdot d\bar{s}$ , where  $s$  is the surface of the sphere  $x^2+y^2+z^2 = a^2$ . [04]

c) If  $\bar{E} = \nabla\phi$ , and  $\nabla^2\phi = -4\pi\rho$ , [04]  
 prove that  $\iint_s \bar{E} \cdot d\bar{s} = -4\pi \iiint_v \rho dv$

Q7) a) Find the value of  $p$  such that the function [04]  
 $f(x) = r^2 \cos 2\Theta + i r^2 \sin p\Theta$  becomes analytical function.

b) Evaluate  $\oint_c \frac{z^2 + \cos^2 z}{(z - \frac{\pi}{4})^3} dz$  [05]

where  $c$  is a circle  $x^2+y^2=1$

c) Find the bilinear transformation which maps  $1, i, -1$  from  $z$  plane into  $i, 0, -i$  from the  $w$  plane [04]

**OR**

Q8) a) Determine the analytic function  $f(z)$  whose real part is [04]

$$U = x^3 - 3xy^2 + 3x^2 - 3y^2 + 1$$

b) Evaluate  $\oint_c \left[ \frac{\sin \pi z^2 + 2z}{(z-1)^2(z-2)} \right] dz$  [05]

where  $c$  is a circle  $x^2+y^2=16$

c) Show that the transformation  $\omega = \sin z$  transforms the straight lines  $x = c$  of  $z$  plane into hyperbolas in the  $\omega$  plane [04]

\*\*\*\*\* END \*\*\*\*\*