

Total No. of Questions—8]

[Total No. of Printed Pages—5

<b>Seat No.</b>	
---------------------	--

**[5252]-140**

**S.E. (Electronics/E&TC) (Second Semester)**

**EXAMINATION, 2017**

**ENGINEERING MATHEMATICS-III**

**(2012 PATTERN)**

**Time : Two Hours**

**Maximum Marks : 50**

**N.B. :—** (i) Attempt Q. 1 or 2, Q. 3 or 4, Q. 5 or 6, Q. 7 or 8.

(ii) Neat diagram must be drawn wherever necessary.

(iii) Figures to the right indicate full marks.

(iv) Use of non-programmable pocket calculator (electronic is allowed).

(v) Assume suitable data, if necessary.

**1. (a) Solve any two :**

**[8]**

(i)  $(D^2 + 2D + 1) y = 2\cos x + 3x + 2$

(ii)  $\frac{d^2y}{dx^2} + y = \operatorname{cosec} x$  (by method of variation of parameter)

(iii)  $x^2 \frac{d^2y}{dx^2} - 2x \frac{dy}{dx} - 4y = x^2 + 2 \log x.$

(b) Find Fourier cosine transform of  $f(x) = \begin{cases} x, & 0 \leq x \leq a \\ 0, & x > a \end{cases}$  **[4]**

**P.T.O.**

Or

2. (a) An electric circuit consists of an inductance 0.1 henry, a resistance (R) of  $20 \Omega$  and a condenser of capacitance (C) of 25 microfarads. If the differential equation of electric circuit

is  $L \frac{d^2q}{dt^2} + R \frac{dq}{dt} + \frac{q}{C} = 0$ , then find the charge 'q' and current 'i' at any time  $t$ , given that at  $t = 0$ ,  $q = 0.05$  coulombs.

$$i = \frac{dq}{dt} = 0 \text{ when } t = 0. \quad [4]$$

- (b) Solve (any one) : [4]

(i) Find  $z$ -transform of  $f(k) = 2^k \cos(3k + 2)$ .

(ii) Find inverse  $z$ -transform of  $\frac{1}{\left(z - \frac{1}{2}\right)\left(z - \frac{1}{3}\right)}$  for  $\frac{1}{3} < |z| < \frac{1}{2}$ .

- (c) Solve : [4]

$$f(k + 1) + \frac{1}{2}f(k) = \left(\frac{1}{2}\right)^k, \quad k \geq 0, \quad f(0) = 0.$$

3. (a) Solve the equation  $\frac{dy}{dx} = \sqrt{x+y}$  using fourth order Runge-Kutta method given  $y(0) = 1$  to find  $y$  at  $x = 0.2$  taking  $h = 0.2$ .

[4]

- (b) Find Lagrange's interpolating polynomial passing through set of points :

$x$	0	1	2
$y$	2	1	4

Hence find  $y$  at  $x = 0.5$  and  $\frac{dy}{dx}$  at  $x = 2$ . [4]

- (c) Find the directional derivative of  $\phi = 4xz^3 - 3x^2y^2z$  at  $(2, -1, 2)$  in the direction of  $2\bar{i} - 3\bar{j} + 6\bar{k}$ . [4]

Or

4. (a) Show that (any one) : [4]

- (i) For scalar functions  $\phi$  &  $\psi$ , show that :

$$\nabla \cdot (\phi \nabla \psi - \psi \nabla \phi) = \phi \nabla^2 \psi - \psi \nabla^2 \phi$$

(ii)  $\nabla \cdot \left( r \nabla \frac{1}{r^3} \right) = \frac{3}{r^4}$

- (b) Show that the vector field  $\bar{F} = (y^2 \cos x + z^2)\bar{i} + 2y \sin x \bar{j} + 2xz \bar{k}$  is irrotational. Find scalar  $\phi$  such that  $\bar{F} = \nabla \phi$ . [4]

- (c) Evaluate  $\int_0^{\pi/2} \frac{\sin x}{x} dx$  by using Simpson's  $\left(\frac{1}{3}\right)^{rd}$  rule by dividing the interval into four parts. Considering the values upto four decimals. [4]

5. (a) Evaluate  $\int_c \bar{F} \cdot d\bar{r}$  for  $\bar{F} = (5xy - 6x^2)\bar{i} + (2y - 4x)\bar{j}$  along the curve  $c : y = x^3$  in XOY plane from  $(1, 1)$  to  $(2, 8)$ . [4]

- (b) Use divergence theorem to evaluate :

$$\iiint_s (y^2 z^2 \bar{i} + z^2 x^2 \bar{j} + x^2 y^2 \bar{k}) \cdot d\bar{S}$$

where S is the upper part of the sphere  $x^2 + y^2 + z^2 = 9$  above XOY plane. [5]

- (c) Using Green's theorem, show that the area bounded by a simple closed curve C is given by :

$$\frac{1}{2} \int_c (x dy - y dx)$$

Hence find area of the ellipse

$$x = a \cos\theta, y = b \sin\theta. \quad [4]$$

Or

6. (a) Find the work done in moving a particle from A(1,0,1) to B(2,1,2) along the straight line AB in the force field  $\bar{F} = x^2\bar{i} + (x-y)\bar{j} + (y+z)\bar{k}$ . [4]

- (b) Evaluate :

$$\iint_s (\nabla \times \bar{F}) \cdot d\bar{S}$$

for the vector field  $\bar{F} = 4y\bar{i} - 4x\bar{j} + 3\bar{k}$  where S is a disc of radius 1 lying on the plane  $z = 1$ . [5]

- (c) Prove that :

$$\iint_s (\phi \nabla \psi - \psi \nabla \phi) \cdot d\bar{S} = \iiint_v (\phi \nabla^2 \psi - \psi \nabla^2 \phi) dV$$

where S is any closed surface enclosing volume V. [4]

7. (a) If  $f(z) = u + iv$  is an analytic function with  $u = \cosh x \cos y$ , express  $f(z)$  in terms of  $z$ . [4]
- (b) Evaluate :

$$\oint_c \frac{2z^2 + z + 5}{\left(z - \frac{3}{2}\right)^2} dz$$

where C is  $\frac{x^2}{4} + \frac{y^2}{9} = 1$ . [4]

- (c) Find the bilinear transformation which maps the points  $0, -i, -1$  from  $z$ -plane into the points  $i, 1, 0$  of the  $w$ -plane. [5]

*Or*

8. (a) If  $f(z) = u + iv = f(re^{i\theta})$  is analytic, show that  $u$  satisfies the Laplace equation  $\frac{\partial^2 u}{\partial r^2} + \frac{1}{r} \frac{\partial u}{\partial r} + \frac{1}{r^2} \frac{\partial^2 u}{\partial \theta^2} = 0$ . [4]
- (b) Evaluate  $\oint_c \frac{e^{2z}}{z(z-1)^2} dz$  over  $c : |z| = 3$ . [4]
- (c) Show that the bilinear transformation  $w = \frac{2z+3}{z-4}$  maps the circle  $x^2 + y^2 - 4x = 0$  into the line  $4u + 3 = 0$ . [5]