

Total No. of Questions—8]

[Total No. of Printed Pages—4+2

Seat No.	
-------------	--

[4956]-108

**F.E. EXAMINATION, 2016**  
**ENGINEERING MATHEMATICS—II**  
**(2012 PATTERN)**

**Time : Two Hours**

**Maximum Marks : 50**

**N.B. :—** (i) Attempt *four* questions : Q. No. 1 or Q. No. 2, Q. No. 3 or Q. No. 4, Q. No. 5 or Q. No. 6, Q. No. 7 or Q. No. 8.

(ii) Neat diagrams must be drawn wherever necessary.

(iii) Figures to the right indicate full marks.

(iv) Use of electronic non-programable calculator is allowed.

(v) Assume suitable data, if necessary.

1. (a) Solve the following differential equations : [8]

(i) 
$$\frac{dy}{dx} = \frac{2x - 3y + 1}{3x + 4y - 5}$$

(ii) 
$$x \cos x \frac{dy}{dx} + (\cos x - x \sin x) y = 1.$$

(b) Assuming that the resistance to movement of a ship through water in the form of  $(a^2 + b^2 v^2)$ , where  $v$  is the velocity,  $a$  and  $b$  are constants, write down the differential equation

P.T.O.

for retardation of ship moving with engine stopped. Prove that the time in which the speed falls to one half its original value  $u$  is given by :

$$\frac{w}{abg} \tan^{-1} \left( \frac{abu}{2a^2 + b^2 u^2} \right)$$

where  $w$  is the weight of the ship. [4]

*Or*

2. (a) Solve : [4]

$$y \log y \, dx + (x - \log y) \, dy = 0$$

(b) Solve the following : [8]

(i) A body of temperature  $80^\circ\text{F}$  is placed in a room of constant temperature  $50^\circ\text{F}$  at time  $t = 0$ . At the end of 5 minutes the body was cooled to a temperature of  $70^\circ\text{F}$ . Find the time at which temperature of the body will be  $60^\circ\text{F}$ .

(ii) A capacitor  $C = 0.01 \text{ F}$  in series with a resistor  $R = 20 \, \Omega$  is charged from a battery 10 Volts. Assuming that initially the capacitor is completely uncharged, determine the charge  $Q(t)$  and current  $I(t)$  in the circuit.

3. (a) Find the Fourier series of : [5]

$$f(x) = x^3, \quad -\pi < x < \pi.$$

(b) Evaluate : [3]

$$\int_0^{\infty} \frac{x^9 (1 - x^5)}{(1 + x)^{25}} dx .$$

(c) Trace the following curve (any one) : [4]

(i)  $x = a(t + \sin t), y = a(1 + \cos t)$

(ii)  $r = a \cos 3\theta.$

Or

4. (a) If [4]

$$I_n = \int_0^{\pi/4} \sec^n \theta d\theta,$$

prove that :

$$I_n = \frac{(\sqrt{2})^{n-2}}{n-1} + \frac{n+2}{n-1} I_{n-2}.$$

(b) If [4]

$$f(x) = \int_2^x (x-t) G(t) dt$$

then show that :

$$\frac{d^2 f}{dx^2} - G(x) = 0.$$

(c) Find the length of arc of an Astroid : [4]

$$x^{\frac{2}{3}} + y^{\frac{2}{3}} = a^{\frac{2}{3}}.$$

5. (a) Show that the plane

$$2x - 2y + z + 12 = 0$$

touches the sphere

$$x^2 + y^2 + z^2 - 2x - 4y + 2z - 3 = 0.$$

Also find the point of contact. [5]

- (b) Find the equation of right circular cone passing through (2, -2, 1) with vertex at origin and axis parallel to the line : [4]

$$\frac{x - 2}{5} = \frac{y - 1}{1} = \frac{z + 2}{1}.$$

- (c) Find the equation of right circular cylinder whose axis is :

$$x = 2y = -z$$

and radius is 4. [4]

*Or*

6. (a) Find the equation of the sphere which has its centre at (2, 3, -1) and touches the line : [5]

$$\frac{x + 1}{-5} = \frac{y - 8}{3} = \frac{z - 4}{4}.$$

- (b) Find the equation of the cone with vertex at (1, 2, -3), semi-vertical angle  $\cos^{-1}\frac{1}{\sqrt{3}}$  and the line :

$$\frac{x - 1}{1} = \frac{y - 2}{2} = \frac{z + 1}{-1}$$

as the axis of the cone. [4]

- (c) Find the equation of right circular cylinder of radius 2 with the axis : [4]

$$\frac{x-1}{2} = \frac{y}{3} = \frac{z-3}{1}.$$

7. Attempt any *two* of the following :

- (a) Evaluate : [6]

$$\iint \frac{1}{x^4 + y^2} dx dy$$

over the region

$$y \geq x^2, x \geq 1.$$

- (b) Evaluate : [7]

$$\iiint \frac{dx dy dz}{\sqrt{1-x^2-y^2-z^2}}$$

taken throughout the volume of the sphere :

$$x^2 + y^2 + z^2 = 1$$

in the positive octant.

- (c) Find the area bounded by the parabola

$$y^2 = 4x$$

and the straight line : [6]

$$2x - 3y + 4 = 0.$$

Or

8. Attempt any *two* of the following :

(a) Evaluate : [6]

$$\int_0^{a/\sqrt{2}} \int_y^{\sqrt{a^2-y^2}} \log_e(x^2 + y^2) dx dy .$$

(b) A rod of length  $l$  is divided into two parts at random. Find average of sum of squares of these parts. Also find mean value of rectangle contained by these two segments. [7]

(c) Find the volume common to the cylinders : [6]

$$x^2 + y^2 = a^2 \text{ and}$$

$$x^2 + z^2 = a^2.$$